

The following Table exhibits a comparative view of the measurements of different parts of the skeletons of the *Cervus Megaceros* in the Museum of the Royal Dublin Society, and in the Royal Museum of the University of Edinburgh, with some parts of the Moose. The measurements of the Edinburgh specimen are taken from Professor Jameson's memoir on *organic remains*, in the Supplement to the Encyclopedia Britannica.

	R. D. Soc.		U. of Edin.		Moose	
	Ft.	In.	Ft.	In.	Ft.	In.
Length of the head, ...	1	8 $\frac{1}{2}$	1	8 $\frac{1}{4}$		
Breadth of the skull between the orbits,	0	10 $\frac{1}{2}$	0	9		
Do. of skull at the occiput,	0	8				
Diameter of the orbit, ...	0	2 $\frac{3}{8}$	0	2 $\frac{1}{2}$		
Distance between infra orbi- tar holes across the skull,	0	7				
Length of alveolar processes of the upper jaw, ...	0	6	0	6		
Length of lower jaw, ...	1	5 $\frac{1}{2}$	0	3 $\frac{1}{2}$		
Diam. of foramen magnum,	0	2				

HORNS.

Distance between the ex- treme tips, measured by the skull,	11	10				
Ditto, in a straight line across,	9	2	6	8	3	7
Length of each horn, ...	5	9	5	1		
Greatest breadth of the palm,	2	10				
Length of the beam,	1	9			0	6 $\frac{1}{2}$
Ditto of brow antler,	0	8 $\frac{5}{4}$				
Ditto of sur-antler,	1	4				
Circumference of the beam at root of brow antler,	1	0 $\frac{5}{4}$			0	7 $\frac{1}{2}$