

- Barton Cliff, Hants. London Clay shells in profusion.
- Binstead, near Ryde, Isle of Wight. In the stone quarries, fresh-water shells, as *Bulimus*, *Helix*, *Phasianella*, *Paludina*, and *Planorbis*.
- Blackdown, near Collumpton, Devon. Numerous silicified shells, of great beauty. *Trigonia*, *Venus*, *Corbula*, *Rostellaria*, &c. &c.
- Bognor Rocks, Sussex. *Vermetus*, *Pectunculus*, *Pinna*, *Murex*, &c.
- Bolland. Numerous shells of the Mountain Limestone.
- Bradford. Numerous Oolitic shells. *Avicula*.
- Bramerton Hill, near Norwich. Shells of the Norfolk Crag.
- Breden, near Derby. *Producta*, *Spirifera*, &c. in Magnesian Limestone.
- Brighton. *Plagiostoma*, *Terebratula*, *Ostrea*, *Pecten*, *Inoceramus*, &c. Many species in the chalk.
- Bromley, Kent. Oyster conglomerate. *Pectunculi*, *Pectens*, &c.
- Brora, Scotland. *Pholadomya*, *Sanguinolaria*, and other shells of the Oolite.
- Calbourn, Isle of Wight. Fresh-water Univalves.
- Cambridge. In the Galt and Chalk-marl, the usual shells.
- Castle Hill, near Newhaven, Sussex. In the Tertiary strata, on the summit of the hill. Numerous *Potamides*, *Cyclades*, and other fresh-water shells. *Ostreæ*, with pebbles.
- Cheltenham. Fine shells of the Oolite and Lias.
- Christchurch, Hants. Abounds in the London Clay shells.
- Chute, near Longleat, Wilts. Immense quantities of Green Sand shells.
- Clayton, near Hurst, Sussex. In Chalk-marl, many rare shells; as, *Dolium nodosum* (*Min. Conch.* tab. 326.).
- Clifton. *Spirifera*, *Producta*, &c.
- Coalbrook Dale. *Euomphalus*, and other Silurian and Carboniferous species.
- Cork. In the vicinity. *Cardium Hibernicum* (*Min. Conch.* tab. 82.), &c.