

NAMES OF FOSSILS.	LOCALITIES.
<p>40. <i>Producta Martini</i>. A fossil of the mountain limestone of Europe.</p>	<p>Windsor, Brookfield, Shubenacadie, East River of Pictou, and Cape Breton.</p>
<p>41. <i>P. concinna</i>, <i>Sow.</i> Or allied Species ; carboniferous limestone of Europe.</p>	<p>Cape Breton.</p>
<p>42. <i>P. Lyelli</i>, <i>De Verneuil</i>. Shell with fine striæ, and with long and slender tubes, the most characteristic fossil of the lower carboniferous formation.</p>	<p>Windsor, Horton Bluff, Shubenacadie, Gay's River, Debert River, Minudie, and Cape Breton.</p>
<p>43. <i>Producta Scotica</i>. Smaller than European, if identical ; carboniferous limestone, Europe.</p>	<p>Windsor, Brookfield, Shubenacadie, East River of Pictou, and Cape Breton.</p>
<p>44. <i>P. Spinosa</i> <i>Sow.</i> <i>Quære.</i> Var. of <i>P. Martini</i>. Carboniferous limestone, Europe.</p>	<p>Cape Breton.</p>
<p>45. <i>P. antiquata</i> de <i>Koninck</i>. Coal-measures, Europe.</p>	<p>Brookfield.</p>
<p>46. <i>Encrinus</i>.</p>	<p>Shubenacadie, and East River, Pictou.</p>
<p>47. <i>Fenestella membranacea</i> ? (<i>Retepora membranacea</i>, <i>Phil.</i>) Carboniferous limestone, Europe.</p>	<p>Windsor, Brookfield, the Shubenacadie, and East River, Pictou.</p>
<p>48. <i>Ceriopa spongites</i>, <i>Goldfuss</i>. (pl. 64.) Eiffel, Silurian, and mountain limestone, Ireland.</p>	<p>Windsor and Brookfield.</p>
<p>49. <i>Favosites ramosa</i> ?</p>	<p>Shubenacadie.</p>
<p>50. <i>Cyathophyllum</i>.</p>	<p>Cape Breton.</p>