CHAPTER XXXVII.—On the different Ages of the Metamorphic Rocks.

Age of each set of metamorphic strata twofold—Test of age by fossils and mineral character not available—Test by superposition ambiguous—Conversion of fossiliferous strata into metamorphic rocks—Limestone and shale of Carrara— Metamorphic strata older than the Cambrian rocks—Others of Lower Silurian origin—Others of the Jurassic and Eocene periods—Why scarcely any of the visible crystalline strata are very modern—Order of succession in metamorphic rocks—Uniformity of mineral character—Why the metamorphic strata are less calcareous than the fossiliferous — Page 611

CHAPTER XXXVIII.-Mineral Veins.

Werner's doctrine, that mineral veins were fissures filled from above—Veins of segregation—Ordinary metalliferous veins or lodes—Their frequent coincidence with faults—Proofs that they originated in fissures in solid rock—Veins shifting other veins—Polishing of their walls or "slicken-sides"—Shells and pebbles in lodes—Evidence of the successive enlargement and reopening of veins—Why some veins alternately swell out and contract—Filling of lodes by sublimation from below—Chemical and electrical action—Relative age of the precious metals—Copper and lead veins in Ireland older than Cornish tin—Lead veins in Lias, Glamorganshire—Gold in Russia, California, and Australia—Connection of hot springs and mineral veins—Concluding remarks - . 618

SUPPLEMENT.

British Pliocene Strata—Proofs from fossil shells of a gradual refrigeration of climate in England, at the successive periods of the Coralline, the Red, and the Norwich Crag—Searles Wood's Monograph on the Crag Mollusca—The Crag Mastodon, a Pliocene species—Different assemblages of fossil Mammalia in the freshwater and drift deposits of the valley of the Thames—Fossil Musk-buffalo in the drift near London and near Berlin – – – – – – Page 685

Where to draw the line between the Miocene and Eocene Tertiary Strata.

Classification of the Miocene and Eocene strata—Where to draw the line between Upper Eocene and Lower Miocene—Reasons for a proposed change of nomenclature—Miocene fossil shells and quadrupeds of the Sewalik or Sub-Himalayan Hills - - 640

Miocene Fuuna of the Sewalik Hills

Denudation of the Wealden-Discovery of the Lower Crag on the summit of the North Downs between Folkestone and Dorking - - - - 645

New Fossil Mammalia from the Purbeck or Upper Oolitic Strata in Dorsetshire.

- Discovery in Dorsetshire of soven or eight new genera of Mammalia in the Purbeck or Upper Oolite strata—First example of a skull of a Mammifer from Secondary Rocks—Insectivorous Marsupials and Placentals and herbivorous Marsupials— Figures and descriptions—Light thrown on the Microlestes or oldest triassic Mammifer—General bearing of the new facts - - - - - - - - - - - - 647
- Discovery of Mammalian Remains in Rocks of high Antiquity in North Carolina, United States - - - 650

Upper Trias of the Euslern Alps.—Recognition of a Marine equivalent of the Upper Trias in the Austrian Alps.—True position of the St. Cassian and Hallstatt Beds.— 800 new species of triassic Mollusca and Radiata—Links thus supplied for connecting the Paleozoic and Neozoic faunas

664 669

645