

- Sections at different points, from Highgate, in Northern Vermont, to Rensselaer County, New York, by Dr. Emmons, 70.
- Cambrian, Table showing correlations of, 44.
- Correlation of, 40.
- Group of, taken across New Hampshire and Vermont, by Prof. C. H. Hitchcock, 20.
- Selwyn, A. R. C., 24, 26, 71, 145.
- Serpulites, in Upper, but not in Middle or Lower Cambrian, 62.
- Species of, in Cambrian, 60.
- dissolutus, 89.
- Shumardia, 153, 157.
- Silurian, Lower (Ordovician), 11, 12.
- Mixing of, with Upper Cambrian, 12.
- Silver Peak, Nevada, 36, 74.
- species from, 38.
- Siphonia, 79.
- Solenopleura, 19, 56, 214.
- common to Lower and Middle Cambrian, 62.
- sp. ? 26.
- Species of, in Cambrian, 61.
- brachymetopa, 215.
- Nana, 27, 48, 214.
- Nana, Explanation of figure of, on plate xxvii, 330.
- (like *S. Nana*), 29.
- Solenosema, 206.
- Spongiae, 50, 72.
- Distribution of species of, in Middle Cambrian, 45.
- Genera and species of, 48.
- Genera and species of, in Cambrian, 62.
- Genera of, in Cambrian, 59.
- Stampede Gap, section near, 33.
- Stenothecca, 128.
- common to Lower, Middle, and Upper Cambrian, 62.
- Species of, in Cambrian, 60.
- Acadica, 53, 128, 129.
- cornucopia, 128, 129.
- elongata, 29, 32, 46, 49, 50, 125, 129.
- elongata, Explanation of figures of, on plate xii, 270.
- pauper, 128, 129.
- rugosa, 26, 27, 29, 46, 49, 53, 126, 128, 129, 139.
- rugosa, Explanation of figures of, on plate xii, 270.
- Straits of Belle Isle, Georgia fauna occurs at the, 29.
- Straparollina, Species of, in Cambrian, 60.
- Strophochetus, 91.
- Species of, in Cambrian, 59.
- ? sp. ?, 38, 45, 91.
- oscellatus, 91.
- Strophomenidae, 52.
- Styliola, 131.
- Swanton, section east of, 18.
- Section east of, by Sir William Logan, 19.
- Section, 24.
- T.
- Taconic, On the use of the name, 65.
- of Emmons considered by Dana as Lower Silurian, 70.
- graptolites, 92.
- Taconic—Continued.
- rocks, Upper division of the, 70.
- series, 66.
- slate, 13.
- system, 65.
- system, Dr. Emmons's section illustrating the, 67.
- system. Position and relation of, by Emmons, 66.
- system, Proposition by Dr. Emmons to divide the, 69.
- Upper, 23.
- Upper, classified by Dr. Emmons as pre-Potsdam, 65.
- Tentaculidæ, 131.
- Tentaculites, 131.
- Tetradium, Palænigma proposed in place of, 223.
- Wrangeli, 223.
- Theca, 131.
- triangularis, 132, 133.
- Topsail Head, Georgia fauna occurs at, 29.
- limestone, correlated by Murray, 49.
- Trails of annelids, in Middle Cambrian, 72.
- Trematis, 102.
- ? pannula, 105.
- punctata, 105.
- Siluriana, 105.
- Trenton limestone, 24.
- Triarthrella, in Upper, but not in Middle or Lower Cambrian, 62.
- Species of, in Cambrian, 61.
- Triarthrus, Becki, 176, 204.
- trilineatus, 203.
- Trichospongia sericea, 78.
- Trilobita, 55.
- Trinucleus concentricus, 152.
- Trinucleus ornatus, 152.
- Triplesia, in Upper, but not in Middle or Lower Cambrian, 62.
- Species of, in Cambrian, 60.
- primordialis, 53, 122.
- Trochocystites Bohemica, 95.
- Trois Pistoles, List of species from, 26.
- section, as described by Sir William Logan, 26.
- Troy, N. Y., 26.
- section, by S. W. Ford, 27.
- U.
- Upper Cambrian, 19, 99.
- Utica shale, 24, 43.
- V.
- Vaginella, 131.
- Vermont, Northern, Potsdam formation not observed in, as it exists in New York, 23.
- section, 28.
- sections, by C. H. Hitchcock, 20.
- Western, sections, 13.
- Von Jhernig, 138.
- W.
- Wasatch section, 38.
- correlated, 44.