FITZINGER, origin of sheep, i. 98; African maned sheep, i. 100.

FITZPATRICK, Mr., potato-grafting, i. 421.

FIXEDNESS of character, conditions of, discussed, ii. 37-39.

FLAX, found in the Swiss lake-dwellings, i. 335; climatal difference in products of, ii. 264.

FIECE, fineness of, in Austrian merinos, ii. 181.

FLEISCHMANN, on German sheep crossed with merinos, ii. 66.

"FLORENTINER-TAUBE," i. 149, 150.

FLOUNDER, ii. 28.

FLOURENS, crossing of wolf and dog, i. 33; prepotency of the jackal over the dog, ii. 43; hybrids of the horse and ass, ii. 43; breeding of monkeys in Europe, ii. 135.

FLOWER-GARDEN, earliest known, in

Europe, ii. 202.

FLOWERS, capricious transmission of colour-varieties in, i. 463, 464; tendency to uniformity in striped, ii. 45; scorching of, dependent on colour, ii. 214; change in, caused by conditions of life, ii. 262; rudimentary, ii. 307; relative position of, to the axis, ii. 337.

FŒTATION, abdominal, ii. 285.

FOLEY, Mr., wild varieties of pears, ii. 248.

FOLIAGE, inherited peculiarities of, i. 385; variegation of, i. 390; bud-variation in, i. 407-409.

Food, influence of, on the pig, i. 76; on cattle, i. 94; excess of, a cause of va-

riability, ii. 244.

FORBES, D., on Chilian sheep, i. 99; on the horses of Spain, Chili, and the Pampas, i. 54.

Formica rufa, ii. 238.

FORTUNE, R., sterility of the sweet potato in China, ii. 153; development of axillary bulbs in the yam, ibid.

Fowl, common, breeds of, i. 236-242; supposed plurality of origin, i. 242; early history of, i. 243-245; causes of production of breeds of, i. 245; origin of, from Gallus bankiva, i. 247-251, 257; feral, notices of, i. 249, 250; reversion and analogous variation in, i. 251-258, ii. 9, 12-14,

341, 343; "cuckoo" sub-breeds of, i. 256; history of, i. 258-260; structural characters of, i. 260-262; sexual peculiarities of, i. 263-270, ii. 50; external differences of, i. 270-273; differences of breeds of, from G. b mkiva, i. 272; osteological characters of, 273-284; effects of disuse of parts in, i. 284-288, ii. 288; feral, i. 199, ii. 7; polydactylism in, i. 458; fertility of, increased by domestication, ii. 90, 145; sterility of, under certain conditions, ii. 144; influence of selection on, ii. 180, 182, 194; evils of close inter-breeding of, ii. 104, 105; crossing of, ii. 73-75; prepotency of transmission in, ii. 42; rudimentary organs in, ii. 306; crossing of non-sitting varieties of, ii. 18, 19; homology of wing and leg feathers in, ii. 315; hybrids of, with pheasants and Gallus sonneratii, ii. 20; black-skinned, ii. 194; black, preyed upon by the osprey in Ireland, ii. 215: five-toed, mentioned by Columella, ii. 424; rumpless, tailed chickens produced by, ii. 4; Dorking, crosses of, ii. 70; form of comb and colour of plumage in, ii. 224; game, crossing of white and black, ii. 70; fivespurred, ii. 386; Spanish, liable to suffer from frost, ii. 296; Polish, peculiarities of skull of, ii. 326-327.

Fox, sterility of, in captivity, ii. 134. Fox, S. BEVAN, races of bees, i. 314.

Fox, W. Darwin, gestation of the dog, i. 31; "Negro" cat, i. 48; reversion of sheep in colour, ii. 3; period of gestation in the pig, i. 77; young of the Himalayan rabbit, i. 114; crossing of wild and domestic turkeys, i. 308; reversion in crossed musk ducks, ii. 14; spontaneous segregation of varieties of geese, ii. 82; effects of close inter-breeding upon bloodhounds, ii. 100; deafness of white cats with blue eyes, ii. 322.

FOXHOUNDS, i. 42, ii. 99.
Fragaria chiloensis, i. 373.
Fragaria collina, i. 373.
Fragaria dioica of Duchesne, i. 375.
Fragaria elatior, i. 373.
Fragaria grandiflora, i. 373.
Fragaria vesca, i. 373.