

- NARVAEZ**, on the cultivation of native plants in Florida, i. 329.
- Nasua**, sterility of, in captivity, ii. 134.
- "**NATAS**," or Niatas, a South American breed of cattle, i. 92-94.
- NATHUSIUS**, H. VON, on striped horses, i. 62; on the pigs of the Swiss lake-dwellings, i. 71; on the races of pigs, i. 68-72; convergence of character in highly-bred pigs, i. 76, ii. 227; causes of changes in the form of the pig's skull, i. 75, 76; changes in breeds of pigs by crossing, i. 82; change of form in the pig, ii. 269; effects of disuse of parts in the pig, ii. 289; period of gestation in the pig, i. 77; appendages to the jaw in pigs, i. 79; on *Sus pliociceps*, i. 72; period of gestation in sheep, i. 101; on Niata cattle, i. 93; on short-horn cattle, ii. 96; on interbreeding, ii. 95; in the sheep, ii. 98; in pigs, ii. 101; unconscious selection in cattle and pigs, ii. 198; variability of highly-selected races, ii. 225.
- NATO**, P., on the Bizzarria orange, i. 417.
- NATURAL** selection, its general principles, i. 2-14.
- NATURE**, sense in which the term is employed, i. 6.
- NAUDIN**, supposed rules of transmission in crossing plants, ii. 43; on the nature of hybrids, ii. 23, 24; essences of the species in hybrids, ii. 382, 395; reversion of hybrids, ii. 9, 23, 24; reversion in flowers by stripes and blotches, ii. 11; hybrids of *Linaria vulgaris* and *purpurea*, ii. 72; pelorism in *Linaria*, ii. 32, 38; crossing of peloric *Linaria* with the normal form, ii. 46; variability in *Datura*, ii. 254; hybrids of *Datura lœvis* and *stramonium*, i. 425; prepotency of transmission of *Datura stramonium* when crossed, ii. 43; on the pollen of *Mirabilis* and of hybrids, i. 415; fertilisation of *Mirabilis*, ii. 356; cultivated Cucurbitaceæ, i. 380, 384, ii. 86; rudimentary tendrils in gourds, ii. 306; dwarf *Cucurbitæ*, ii. 324; relation between the size and number of the fruit in *Cucurbita pepo*, ii. 336; analogous variation in *Cucurbitæ*, ii. 341; acclimatisation of Cucurbitaceæ, ii. 304; production of fruit by sterile hybrid Cucurbitaceæ, ii. 156; on the melon, i. 383, ii. 86, 264; incapacity of the cucumber to cross with other species, i. 382.
- NECTARINE**, i. 357-365; derived from the peach, i. 357, 360-364; hybrids of, i. 360; persistency of characters in seedling, *ibid.*; origin of, i. 361; produced on peach-trees, i. 361, 362; producing peaches, i. 361; variation in, i. 363, 364; bud-variation in, i. 398; glands in the leaves of the, ii. 217; analogous variation in, ii. 341.
- NECTARY**, variations of, in pansies, i. 393.
- NEES**, on changes in the odour of plants, ii. 264.
- "**NEGRO**" cat, i. 48.
- NEGROES**, polydactylism in, i. 548; selection of cattle practised by, ii. 191.
- NEOLITHIC** period, domestication of *Bos longifrons* and *primigenius* in the, i. 84; cattle of the, distinct from the original species, i. 90; domestic goat in the, i. 105; cereals of the, i. 334.
- NERVE**, optic, atrophy of the, ii. 287.
- NEUBERT**, potato-grafting, i. 422.
- NEUMEISTER**, on the Dutch and German pouter pigeons, i. 145; on the Jacobin pigeon, i. 162; duplication of the middle flight feather in pigeons, i. 167; on a peculiarly coloured breed of pigeons, "Staarhalsige Taube," i. 169; fertility of hybrid pigeons, i. 201; mongrels of the trumpeter pigeon, ii. 42; period of perfect plumage in pigeons, ii. 53; advantage of crossing pigeons, ii. 106.
- NEURALGIA**, hereditary, ii. 55.
- NEW ZEALAND**, feral cats of, i. 49; cultivated plants of, i. 328.
- NEWFOUNDLAND** dog, modification of, in England, i. 44.
- NEWMAN**, E., sterility of Sphingidæ under certain conditions, ii. 141.
- NEWPORT**, G., non-copulation of *Vanessa* in confinement, ii. 141; fertilisation of the ovule in batrachia, ii. 356.