

PHTHISIS, affection of the fingers in, ii. 326.

PHYLOXERA, i. 354.

PICKERING, Dr., on the grunting voice of humped cattle, i. 83; occurrence of the head of a fowl in an ancient Egyptian procession, i. 258; seeding of ordinarily seedless fruits, ii. 152; extinction of ancient Egyptian breeds of sheep and oxen, ii. 421; on an ancient Peruvian gourd, ii. 425.

PICOTEES, effect of conditions of life on, ii. 262.

PICTET, A., oriental names of the pigeon, i. 215.

PICTET, Prof., origin of the dog, i. 15; on fossil oxen, i. 85.

PIEBALDS, probably due to reversion, ii. 11.

PIÉTREMENT, M., on the ribs of horses, i. 52.

PIGEAUX, hybrids of the hare and rabbit, ii. 135.

PIGEON à cravate, i. 156.

PIGEON bagadais, i. 149, 150.

PIGEON coquille, i. 163.

PIGEON cygne, i. 150.

PIGEON heurté, i. 163.

PIGEON pattu plongeur, i. 164.

PIGEON polonais, i. 151.

PIGEON romain, i. 149, 150.

PIGEON tambour, i. 162.

PIGEON turc, i. 146.

PIGEONS, origin of, i. 137, 140, 189-214; classified table of breeds of, i. 142; pouter, i. 143-145; carrier, i. 146-149; runt, i. 149-151; barbs, i. 151-153; fantail, i. 153-155; turbit and owl, i. 156; tumbler, i. 156-161; Indian frill-back, i. 161; Jacobin, *ibid.*; trumpeter, i. 162; other breeds of, i. 163-165; differences of, equal to generic, i. 166; individual variations of, i. 167-169; variability of peculiarities characteristic of breeds in, i. 169; sexual variability in, i. 170, 171; osteology of, i. 171-177; correlation of growth in i. 177-180, ii. 313; young of some varieties naked when hatched, i. 179, ii. 326; effects of disuse in, i. 180-187; settling and roosting in trees, i. 190; floating in the Nile to drink, *ibid.*; dovecot, i. 194, 195; arguments for

unity of origin of, i. 197-214; feral, in various places, i. 199, ii. 7; unity of coloration in, i. 204-206; reversion of mongrel, to coloration of *C. livia*, i. 207-214; history of the cultivation of, i. 215-217; history of the principal races of, i. 217-222; mode of production of races of, i. 222-235; reversion in, ii. 22; by age, ii. 14; produced by crossing in, ii. 14, 22; prepotency of transmission of characters in breeds of, ii. 41, 42; sexual differences in some varieties of, ii. 50; period of perfect plumage in, ii. 53; effect of segregation on, ii. 63; preferent pairing of, within the same breed, ii. 81; fertility of, increased by domestication, ii. 90, 138; effects of inter-breeding and necessity of crossing, ii. 106; indifference of, to change of climate, ii. 144; selection of, ii. 79, 183, 188; among the Romans, ii. 187; unconscious selection of, ii. 196, 198; facility of selection of, ii. 220; white, liable to the attacks of hawks, ii. 215; effects of disuse of parts in, ii. 288; fed upon meat, ii. 294; effect of first male upon the subsequent progeny of the female, i. 437; homology of the leg and wing feathers in, ii. 315; union of two outer toes in feather-legged, *ibid.*; correlation of beak, limbs, tongue, and nostrils, ii. 316; analogous variation in, ii. 341, 342; permanence of breeds of, ii. 425.

PIGS, of Swiss lake-dwellings, i. 71; types of, derived from *Sus scrofa* and *Sus indicus*, i. 68-70; Japanese (*Sus pliciceps*, Gray), figured, i. 72; of Pacific Islands, i. 73, ii. 64; modifications of skull in, i. 72-75; length of intestines in, i. 77, ii. 293; period of gestation of, i. 77; number of vertebrae and ribs in, *ibid.*; anomalous forms, i. 78, 79; development of tusks and bristles in, i. 79; striped young of, i. 80; reversion of feral, to wild type, i. 80, 81, ii. 7, 22; production and changes of breeds of, by inter-crossing, i. 82; effects produced by the first male upon the subsequent progeny of the female, i. 436; pedigrees of, i. 447;