

- RAM, goat-like, from the Cape of Good Hope, ii. 41.
- RAMU, M., on appendages to throat of goat, i. 106.
- RANCHIN, heredity of diseases, i. 451.
- RANGE of gallinaceous birds on the Himalaya, i. 249.
- RANKE, on the effects of use and disuse of organs, ii. 285.
- Ranunculus ficaria*, ii. 154.
- Ranunculus repens*, ii. 152.
- RAPE, i. 344.
- Raphanus caudatus*, i. 345.
- Raphanus raphanistrum*, i. 345.
- Raphanus sativus*, ii. 335.
- RASPBERRY, yellow-fruited, ii. 216.
- RATTLESNAKE, experiments with poison of the, ii. 279.
- RAVEN, stomach of, affected by vegetable diet, ii. 292.
- RAWSON, A., self-impotence in hybrids of *Gladiolus*, ii. 121, 122.
- RÉ, COMTE, on the assumption of a yellow colour by all varieties of maize, i. 339.
- RÉAUMUR, effect of confinement upon the cock, ii. 27; fertility of fowls in most climates, ii. 144.
- REED, Mr., atrophy of the limbs of rabbits, consequent on the destruction of their nerves, ii. 287.
- REGENERATION of amputated parts in man, i. 459; in the human embryo, ii. 357; in the lower vertebrata, insects, and myriapoda, ii. 358.
- RE-GROWTH of amputated joints, ii. 357.
- REGNIER, early cultivation of the cabbage by the Celts, i. 342; selection practised by the Celts, ii. 187.
- REINDEER, individuals recognised by the Laplanders, ii. 238.
- REISSEK, experiments in crossing *Cytisus purpureus* and *laburnum*, i. 416; modification of a *Thesium* by *Æcidium*, ii. 274.
- RELATIONS, characters of, reproduced in children, ii. 7.
- RENGGER, occurrence of jaguars with crooked legs in Paraguay, i. 17; naked dogs of Paraguay, i. 24, 32, ii. 70, 80; feral dogs of La Plata, i. 28; on the aguara, i. 27; cats of Paraguay, i. 48, ii. 63, 133; dogs of Paraguay, ii. 64; feral pigs of Buenos Ayres, i. 80; on the refusal of wild animals to breed in captivity, ii. 132; on *Dicotyles labiatus*, ii. 133; sterility of plantigrade carnivora in captivity, ii. 134; on *Cavia aperca*, *ibid.*; sterility of *Cebus azaræ* in captivity, ii. 136; abortions produced by wild animals in captivity, ii. 142.
- REPRODUCTION, sexual and asexual, contrasted, ii. 351; unity of forms of, ii. 378; antagonism of, to growth, ii. 379.
- Reseda odorata*, self-sterility of, ii. 118, 223.
- RETINITIS, pigmentary, in deaf-mutes, ii. 322.
- REUTER, Herr, potato-grafting, i. 422.
- REVERSION, ii. 1, 2, 368, 369, 390, 393-399; in pigeons, ii. 2; in cattle, ii. 3; in sheep, *ibid.*; in fowls, ii. 4; in the heartsease, *ibid.*; in vegetables, *ibid.*; in feral animals and plants, ii. 5-7; to characters derived from a previous cross in man, dogs, pigeons, pigs, and fowls, ii. 7-9; in hybrids, ii. 9; by bud-propagation in plants, ii. 10-13; by age in fowls, cattle, &c., ii. 12, 13; partial, from an injury, ii. 12; caused by crossing, ii. 13-25; explained by latent characters, ii. 25-31; producing monstrosities, ii. 31; producing peloric flowers, ii. 32-34; of feral pigs to the wild type, i. 81, 82; of supposed feral rabbits to the wild type, i. 108, 116, 118; of pigeons, in coloration, when crossed, i. 206-213; in fowls, i. 251-258; in the silkworm, i. 318; in the pansy, i. 393; in a pelargonium, i. 403; in Chrysanthemums, i. 404; of varieties of the China rose in St. Domingo, i. 406; by buds in pinks and carnations, i. 407; of lacinated varieties of trees to the normal form, i. 408; in variegated leaves of plants, i. 409; in tulips, i. 412; of suckers of the seedless barberry to the common form, i. 410; by buds in hybrids of *Tropæolum*, i. 425; in plants, i. 442; of crossed peloric snapdragons, ii. 46; analogous variations due to, ii. 340-344.