

- i. 35; deficiency of, in naked Turkish dogs, i. 36; period of appearance of, in breeds of dogs, *ibid.*; precocity of, in highly-bred animals, ii. 313; correlation of, with hair, ii. 319; double row of, with redundant hair, in Julia Pastrana, ii. 321; affected in form by hereditary syphilis and by pulmonary tubercle, ii. 325; developed on the palate, ii. 385.
- TIGETMEIER, Mr.**, on a cat with monstrous teeth, i. 50; on a swift-like pigeon, i. 165; on sexual colours, i. 170; naked young of some pigeons, i. 179; fertility of hybrid pigeons, i. 201; on white pigeons, ii. 215; reversion in crossed breeds of fowls, i. 251-256; chicks of the white silk-fowl, i. 261; development of the cranial protuberance in Polish fowls, i. 262; on the skull in the Polish fowl, i. 268, 275; on the intelligence of Polish fowls, i. 276; correlation of the cranial protuberance and crest in Polish fowls, i. 289; development of the web in the feet of Polish fowls, i. 272; early development of several peculiarities in Spanish cocks, i. 263; on the comb in Spanish fowls, i. 266; on the Spanish fowl, ii. 296; varieties of game-fowls, i. 264; pedigrees of game-fowls, i. 447; assumption of female plumage by a game-cock, i. 265; natural selection in the game-cock, ii. 210; pugnacity of game-hens, i. 268; length of the middle toe in Cochin fowls, i. 272; origin of the Sebright bantam, ii. 29; differences in the size of fowls, i. 270; effect of crossing in fowls, *ibid.*; ii. 74; effects of interbreeding in fowls, ii. 105, 106; incubation by mongrels of non-sitting races of fowls, ii. 18; inverse correlation of crest and comb in fowls, i. 288; occurrence of pencilled feathers in fowls, ii. 14; on a variety of the goose from Sebastopol, i. 304; on the fertility of the peahen, ii. 91; on the intercrossing of bees, ii. 107.
- TEMMINCK**, origin of domestic cats, i. 46; origin of domestic pigeons, i. 189; on *Columba guinea*, i. 192; on *Columba leucocephala*, *ibid.*; asserted reluctance of some breeds of pigeons to cross, i. 202; sterility of hybrid turtle-doves, i. 203; variations of *Gallus bankiva*, i. 247; on a buff-coloured breed of turkeys, i. 309; number of eggs laid by the peahen, ii. 91; breeding of guans in captivity, ii. 139; behaviour of grouse in captivity, *ibid.*; sterility of the partridge in captivity, *ibid.*
- TENDRILS** in Cucurbitaceæ, i. 381, ii. 306.
- TENNENT, Sir J. E.**, on the goose, i. 302; on the growth of the apple in Ceylon, ii. 266; on the Jaffna sheep, ii. 292.
- Teredo*, fertilisation in, ii. 356.
- TERRIERS**, wry-legged, ii. 232; white, subject to distemper, ii. 330.
- TESCHEMACHER**, on a husked form of maize, i. 339.
- TESSIER**, on the period of gestation of the dog, i. 30; of the pig, i. 77; in cattle, i. 90; experiments on change of soil, ii. 129.
- Tetrao*, breeding of species of, in captivity, ii. 139.
- Tetrapteryx paradisea*, ii. 140.
- Teucrium campanulatum*, pelorism in, ii. 337.
- TEXAS**, feral cattle in, i. 89.
- THEOGNIS**, his notice of the domestic fowl, i. 258.
- THEOPHRASTUS**, his notice of the peach, ii. 299.
- Thesium*, ii. 274.
- THOMPSON, Mr.**, on the peach and nectarine, i. 363; on the varieties of the apricot, i. 365; classification of varieties of cherries, i. 369; on the "Sister ribston-pippin," i. 372; on the varieties of the gooseberry, i. 376, 377.
- THOMPSON, WILLIAM**, on the pigeons of Ilay, i. 193; feral pigeons in Scotland, i. 199; colour of the bill and legs in geese, i. 304; breeding of *Tetrao scotius* in captivity, ii. 139; destruction of black fowls by the osprey, ii. 215.
- THORN**, grafting of early and late, i. 386; Glastonbury, i. 387.
- THORNS**, reconversion of, into branches, in pear trees, ii. 310