

- tilus, i. 326; ditto in ammonites, i. 351; ditto in belemnites, i. 381.
- Hyanas, bones collected by, in caverns, i. 94.
- Hylæosaurus discovered by Mr. Mantell, i. 241; peculiar character of, i. 241.
- Hythe, large hamite found at, i. 368.
- Icthyodorulites, or fossil spines, i. 288; uses and variety of, i. 290.
- Ichthyosaurus, geological extent and chief localities of, i. 168; curious structure of, i. 169; number of species, i. 169; head partaking of the character of crocodiles and lizards, i. 171; jaw, length of, i. 171; teeth, character and number of, i. 171; how differing from crocodiles, i. 171; contrivances for replacing, i. 172; eyes, magnitude of, i. 173; eyes, microscopic and telescopic properties of, i. 173; eyes, bony sclerotic of, i. 174, 175; jaws, composed of many thin plates, i. 175; jaw, lower, contrivances in, i. 176, 177; vertebræ, number of, i. 177; vertebræ constructed like those of fishes, i. 178; ribs structure of, and to what purpose subservient, i. 180; sternum like that of ornithorhynchus, i. 181; paddles, anterior, like those of whales; posterior, like those of ornithorhynchus, i. 182, 183; concluding remarks upon, i. 185, 186; intestinal structure of, i. 187; skeleton of, containing coprolite, i. 191; small intestines spiral, like that of sharks and rays, i. 193; final cause of spiral intestinal structure, i. 195; skin of preserved, ii. 22; mechanism of atlas and cervical vertebræ of, ii. 24—26.
- Igneous rocks, various phenomena of, ii. 5—9.
- Iguana, modern, habits of, i. 242; den- tition of, i. 248.
- Iguanodon, discovered by Mr. Mantell, i. 240; remains of, where found, i. 241; a gigantic herbivorous reptile, i. 241; teeth like those of the modern iguana, i. 241; the largest of known fossil reptiles, i. 241, 244; climate indicated by remains of, i. 243; teeth, peculiar character of, i. 245—248; bony horn on the nose of, i. 244; food of, probably tough vegetables, i. 246.
- Indusiæ, fossil, in fresh water formation of Auvergne, i. 118.
- Infusoria, Ehrenberg's observations on, i. 446—448; number of species described, i. 446; their powers of reproduction, i. 446; their manners of increase, i. 447; universal diffusion of, i. 448; known extent of living and fossil species, i. 610—612.
- Injection of igneous rocks at various periods, ii. 6.
- Ink bags, recent and fossil of loligo, i. 303.
- Insects, fossil in carboniferous strata, 409; wing covers of, at Stonesfield, i. 411; Count Munster's collection of from Solenhofen, i. 411; many fossil genera in tertiary strata, i. 412; found at Aix by Mr. Murchison and Mr. Lyell, i. 610.
- Iron, ore abundant in coal formation, i. 65; quantity of, annually made in England and Wales, i. 530; secreted by living infusoria, i. 611.
- Isle of Wight, iguanodon found in, i. 242.
- Jæger, Professor, has found coprolites in Wirtemberg, i. 190; his work on fossil plants, i. 490.
- Jardine, Sir W., fossil footmarks found by, i. 259.
- Juli, supposed fir cones, are coprolites, i. 198.
- Kaup, Professor, notice on the footsteps of chirotherium i. 264; his description of fossil mammalia at Epplesheim, i. 91; his description of dinotherium, i. 136, 603.
- Kersten, Professor, his discovery of pyrogenous crystals of felspar, i. 598.
- Kepler, his prayer, i. 10.
- Killery, cemetery in a sand bank at, i. 104.
- King, Captain, animal of spirula found by, i. 362; serolis found by, i. 392.
- König, Mr., his account of human skeleton in British Museum, from Guadeloupe, i. 104.
- LAMANTIN, remains of, in pliocene periods, indicate a warm climate, i. 92.
- Lamarck, his two sections of trachelipods, i. 298; on geological effects of microscopic shells, i. 385; his notice of suckers of encrinus, i. 440; theory of transmutation associated with development by, i. 585.
- Lartet, M., animals at Simorre described by, i. 601.