

contain fossils materially different from those in the under. The clusters of small oyster-shells, and the stems of the pentacrinus, lie near together, and not many other fossils occur toward the bottom of the rock. Both these sorts are found about the clay which lies beneath the stone; and in a detached piece of the Cornbrash Stratum, (not ten acres,) near Pipe-house, south of Bath, the encrini are found above the springs, in the brownish clay turned up by the plough.

---

*ORGANIZED FOSSILS.*

FIG.

- |   |   |
|---|---|
| 1 <i>Natica</i> ?   | Road, Sleaford. Wick Farm.  |
| 2 <i>Ammonites</i> discus                                   | Closworth. Road. S. W. of Wincanton. Chillington.   |
| 3 <i>Modiola</i>  | Closworth. Wick Farm. Holt.   |
| 4 <i>Trigonia</i> costata                                   | North side of Wincanton. Wick Farm.   |
| 5 <i>Venus</i> <i>Linn.</i>                                 | Trowle. Sheldon. S. W. of Wincanton. Norton.  |
| 6 <i>Cardium</i>  | Road. Elmcross. Wick Farm. Sleaford. Woodford. Near Peterborough. Near Stilton.   |
| 7 <i>Unio</i> ?   | North Cheriton. Road. Draycote. Maisey Hampton. Sleaford. S. W. of Tellisford. Sattyford. S. W. of Wincanton.   |
| 8 <i>Avicula</i> echinata                                   | Closworth. North Cheriton. Lullington. Trowle. Sheldon. Draycot. Norton. Stony Stratford. S. W. of Tellisford. North side of Wincanton. S. W. of Wincanton. |
| 9 <i>Terebratula</i> digona ( <i>var.</i> gibbosa, rotunda) | Closworth. Redlynch. Trowle. Wick Farm. Sheldon. Latton. Woodford.  |