

- Crich Hill, Derbyshire. The usual shells of the *Mountain Limestone*.
- Cuckfield, Sussex. In the Sandstone and Grit, fresh-water shells of the Wealden.
- Dowlands, near Lyme Regis. Many fine shells. *Pachymya gigas* (*Min. Conch.* tab. 504.).
- Dudley. Profusion of shells of the Silurian strata.
- Dundry, near Bristol. Beautiful shells in the Inferior Oolite. *Trochus*, *Cardita*, &c.
- Earlstone, Wilts. Many shells of the Green Sand.
- Faringdon, Berks. The usual shells of the Oolite in the Coral Rag, &c.; and of the Green Sand, in the Gravel-pits.
- Folkstone, Kent. *Inoceramus*, *Arca*, *Rostellaria*, *Dentalium*, &c., in the Galt. In the Green Sand, a large *Gryphæa*, *Ostrea*, &c.
- Gravesend. Beautiful shells of the White Chalk.
- Hampstead Cliff, Isle of Wight. Freshwater Tertiary shells.
- Hampton Quarry, near Bath. Abounds in Oolitic shells.
- Hartwell, Bucks. On the estate of Dr. Lee, beautiful shells of the Kimmeridge Clay.
- Harwich Cliff, Essex. The Crag shells. *Voluta Lamberti* (*Min. Conch.* tab. 129.).
- Hastings, Sussex. Fresh-water shells of the Wealden.
- Headon Hill, Isle of Wight. Fresh-water Tertiary shells in profusion.
- Heddington. Oysters in Kimmeridge Clay (*Ostrea deltoidea*). *Perna*, *Gervillia*, *Trigonia*, &c.
- Highworth, Wilts. Very fine *Trigoniæ*, and other Oolitic shells, in the stone-quarries.
- Hollington, near Hastings. Fresh-water bivalves in limestone.
- Holywell, near Ipswich. Shells of the Crag, abundantly.
- Hordwell Cliff, Hants. The usual tertiary shells of the *Eocene* deposits, in immense quantity, variety, and perfection.
- Horningsham, near Frome, Wilts. *Terebratula*, *Pecten*, &c. in great numbers.