

Prematurity has often been recognized as evidence of low development and low rank; and the following is the explanation of it.

When an animal has reached the condition required for locomotion and for the care of itself, it has already the essential faculties of an adult; and although these faculties of locomotion and self-feeding are of comparatively low grade, the animal possessing them is approximately mature in its cephalic forces, and afterwards rises but little with growth. Prematurity hence involves inferiority. The pupa-state of an Insect is a means of higher development the more perfect its inactivity. For this complete rest allows all the forces of the individual to be concentrated on the internal processes, and favors, therefore, that cephalic growth which makes a special demand on these forces; while in an *active* pupa (or rather the larve that passes through no pupa-state), activity, whether that of locomotion, or of digestion, constantly exhausts force; and only the balance, not thus run away with, goes towards the maturing process. With such an open outlet of force, the animal may mature physically, that is, grow and perfect its outer structure; but cephalically, or, in all those points of structure, as well as psychical powers, that are connected with superior cephalic development, it makes little advance.

Hence, (a) those insects whose larves are essentially like the adults and undergo no metamorphosis are inferior in type,—as generally so recognized.

Again, (b) those Insects (as most Hymenopterous) whose larves are footless grubs are superior in type to those (as the Lepidopterous) whose larves are most highly developed and active.

Viewed on the ascending grade, this method is the *permativative*.

13. *Gemmative*.—Exhibited in multiplication by buds. Budding may produce—

a. Perfect individuals, capable of egg-production.

b. Individuals capable only of budding, and giving origin to a perfect egg-producing individual as the last of a series of buddings.

c. Caducous, or persistent buds; the *latter* leading to compound forms, either branching, lamellar, or massive.

This power of reproduction by buds occurs in many Worms, both superior and inferior; in Bryozoan and many Ascidian Mollusks; in Polyps and many other Radiates. The production of persistent buds is the lowest grade, and is common in the budding Mollusks and Radiates, but not the Articulates. Among budding Articulates, case *b* appears to be of lower grade than case *a*.

This method is allied to the *multiplicative*, p. 325. It is also *phytozoic* (p. 327), or a plant-like feature in animal life.