

- Columbia, ii. 189; crossed with wild banteng in Java, ii. 191; with reversed hair in Banda Oriental, ii. 190; selection of trifling characters in, ii. 193; fashion in, ii. 195; similarity of best races of, ii. 227; unconscious selection in, ii. 198; effects of natural selection on anomalous breeds of, ii. 211, 212; light-coloured, attacked by flies, ii. 214, 330; Jersey, rapid improvement of, ii. 220; effects of disuse of parts in, ii. 289; rudimentary horns in, ii. 306; supposed influence of humidity on the hair of, ii. 318; white spots of, liable to disease, ii. 331; supposed analogous variation in, ii. 341; displacement of long-horned by short-horned, ii. 421.
- CAULIFLOWER, i. 341; free-seeding of, in India, ii. 301; rudimentary flowers in, ii. 306.
- CAVALIER pigeon, ii. 75.
- Cavia aperea*, ii. 134.
- CAY (*Cebus azaræ*), sterility of, in confinement, ii. 136.
- Cebus azaræ*, ii. 136.
- Cecidomyia*, larval development of, ii. 273, 353, 363; and *Misocampus*, i. 5.
- CEDARS of Lebanon and Atlas, i. 387.
- CELERY, turnip-rooted, i. 344; run wild, ii. 7.
- CELL-THEORY, ii. 366.
- Celosia cristata*, i. 389.
- CELSUS, on the selection of seed-corn, i. 336, ii. 187.
- CELTS, early cultivation of the cabbage by the, i. 342; selection of cattle and horses by the, ii. 187.
- Cenchrus*, seeds of a, used as food, i. 326.
- Centaurea cyanus*, bud-variation in, i. 404.
- CEPHALOPODA, spermatophores of, ii. 379.
- Cerasus padus*, yellow-fruited, i. 462.
- Cercoleptes*, sterility of, in captivity, ii. 134.
- Cercopithecus*, breeding of a species of, in captivity, ii. 136.
- CEREALS, i. 329, 330; of the Neolithic period in Switzerland, i. 335; adaptation of, to soils, ii. 295.
- Cereus*, ii. 11.
- Cereus speciosissimus* and *phyllanthus*, reversion in hybrids of, i. 425.
- Cervus canadensis*, ii. 141.
- Cervus dama*, ii. 99.
- CETACEA, correlation of dermal system and teeth in the, ii. 321.
- CEYLON, cats of, i. 48; pigeon-fancying in, i. 216.
- CHAMISSO, on seeding bread-fruit, ii. 152.
- CHANNEL Islands, breeds of cattle in, i. 83.
- CHAPMAN, Professor, peach-trees producing nectarines, i. 362.
- CHAPUIS, F., sexual peculiarities in pigeons, i. 170; effect produced by first male upon the subsequent progeny of the female, i. 437; sterility of the union of some pigeons, ii. 146.
- CHARACTERS, fixity of, ii. 225; latent, ii. 25-31, 393, 394; continued divergence of, ii. 227; antagonistic, ii. 395.
- CHARDIN, abundance of pigeons in Persia, i. 215.
- CHARLEMAGNE, orders as to the selection of stallions, ii. 187.
- CHARTLEY, wild cattle of, i. 88.
- CHATÉ, reversion of the upper seeds in the pods of stocks, ii. 340.
- CHAUNDY, Mr., crossed varieties of cabbage, ii. 110.
- CHEETAH, general sterility of, in captivity, ii. 133.
- Cheiranthus cheiri*, i. 407.
- CHERRIES, i. 368, 369; bud-variation in, i. 399; white Tartarian, ii. 215; variety of, with curled petals, ii. 218; period of vegetation of, changed by forcing, ii. 301.
- CHEVREUL, on crossing fruit-trees, ii. 110.
- CHICKENS, differences in characters of, i. 261, 262; white, liable to gapes, ii. 213, 330.
- CHIGOE, ii. 265.
- CHILE, sheep of, i. 99.
- CHILLINGHAM cattle, identical with *Bos primigenius*, i. 84; characters of, i. 87.
- CHILOE, half-castes of, ii. 21.
- CHINA, cats of, with drooping ears, i. 49; horses of, i. 56; striped ponies of, i. 62; asses of, i. 66; notice of rabbits in, by Confucius, i. 107; breeds