

- Saarbrück, fishes found at, i. 266, 278.
- Sabrina island, rise and destruction of, ii. 8.
- Sacred history, consistency of geological discoveries with, i. 8.
- Salado, river, megatherium in bed of, i. 143, 161.
- Salamander, fossil at Oeningen, i. 514; from Japan, alive at Leyden, i. 514.
- Salt, found in secondary and tertiary strata, i. 71.
- Sand, effects of wind in forming strata of, i. 127.
- Sapey Brook, concretions in, mistaken for footsteps, i. 261.
- Saurians, character of in secondary strata, i. 74; sudden death and burial in lias clay, i. 124; fossil history and relations of, i. 165—168; in what formations found, i. 166, 168; amphibious, allied to crocodiles, i. 249; gigantic terrestrial, i. 234; flying, 221; marine, i. 168, 202.
- Scaphite, character and extent of, i. 368.
- Schlotheim, his early arrangement of fossil plants, i. 456.
- Schmerling, M., bones found by, in caves near Liege, i. 602.
- Sciences, geology essential to advancement of, i. 7.
- Scorpions, fossil in coal formation, i. 406; indicate a warm climate, i. 408; fossil, description of, i. 407; eyes and skin, preserved, i. 407, 408; hairs preserved, i. 411.
- Scrope, Mr. Poulett, his panoramic views of Auvergne, ii. 8; on ripple marks and tracks of animals in oolitic strata, i. 260; ii. 42.
- Sea, early history of, illustrated by fossil eyes, i. 401; crowded with animal life, i. 293.
- Secondary strata, history of, i. 67; adaptation of to human uses, i. 68; materials of, whence derived, i. 68; nature of materials, i. 69; advantageous disposition of, i. 70.
- Secondary formations, leading character of their fossil vegetables, i. 453.
- Sedgwick, Professor, on the kind of information to be looked for in the Bible, i. 34, 594; his discovery of fossil fishes, i. 277.
- Segregation, theory of veins filled by, i. 551.
- Sellow, M., his use of Chinese method of boring wells, i. 568.
- Sepiostaire, its analogies to belemnite, i. 378.
- Serolis, allied to trilobites, i. 392, 394.
- Serpent, fossil, in Auvergne, i. 618.
- Serpentine, veins and masses of, ii. 5.
- Serpulæ, attached to belemnites, i. 377.
- Sharks, antiquity of family of, i. 286; extinct species, numerous, i. 286; fossil teeth of, i. 286; fossil spines, or ichthyodorulites, i. 287; three sub-families of, i. 287; teeth in early families obtuse, i. 289; peculiar form of tail, 279.
- Sheerness, Artesian well at, i. 563.
- Shells, number of in tertiary strata, i. 79; vast accumulation of, in many strata, i. 116; turbinated, formed by animals of higher order than bivalves, i. 296; fossil univalve and bivalve, i. 295, 296; bored by carnivorous trachelipods, i. 299; specific gravity of, i. 302; bivalve, constructed by conchifers, i. 296; proofs of design in fossil chambered, i. 310; conclusions from chambered species, i. 380; foraminated polythalamous, i. 381; microscopic, quantity of, in certain strata, i. 117; minute multilocular, i. 382.
- Sheppey, fossil emys at, i. 258; fossil crocodile at, i. 258; fishes in London clay at, i. 285; fossil fruits found at, i. 507, 519.
- Shrewsbury, freshwater limestone in coal formation near, i. 599.
- Sickler, Dr., letter on footsteps at Hessberg, i. 265.
- Siebold, Dr., salamander brought from Japan by, i. 514; silicified buprestis in collection of, ii. 78.
- Sienite, veins and overlying masses of, ii. 5.
- Sigillaria, among the largest and tallest plants of the coal formation, i. 469; stems occasionally found erect, i. 470, 471; stem occasionally divided at the summit, i. 472; character and relations of, i. 472, 473, 474, 618; scars on bark in vertical rows, i. 473; number of species, i. 473.
- Silex, secreted by living infusoria, i. 611, 612.
- Silistria, sturgeons in the Danube near, i. 279.
- Silliman, Professor, his interpretation of the word beginning, and of the days of the Mosaic creation, i. 18.
- Silurus, spine of, i. 290.
- Silurian system, its geological place, and history of its establishment, i. 527; recognition of, on the continent, i. 528; divisions of, ii. 104.